

Colonoscopy Basics

What is a colonoscopy?



An endoscopic procedure which uses a flexible tube with a camera and a light source (colonoscope) to look directly at the inner lining of the large intestine (colon)


When to get a colonoscopy?

- Colon cancer screening
- History of precancerous/adenomatous polyps
- Changes in bowel habits
- Iron deficiency anemia
- Rectal bleeding

Before your procedure

- Obtain the bowel prep from your pharmacy.
- **STOP iron** supplements **one week** before.
- **STOP** glucagon-like peptide agonist (GLP-1) like **semaglutide** (Ozempic, Wegovy), **dulaglutide** (Trulicity), **exenatide** (Byetta or Bydureon), or **liraglutide** (Victoza, Saxenda) **one week** before.
- Let your doctor know if you take a **blood thinner** like coumadin (Warfarin), clopidogrel (Plavix), dabigatran (Pradaxa), rivaroxaban (Xarelto), or apixaban (Eliquis) so they can tell you when to hold them.

	Starting 3 Days Before Colonoscopy	2 Days Before Colonoscopy	The Day before Colonoscopy	On Colonoscopy Day
Diet	Low Residue Diet	Low Residue Diet	Clear Liquid Diet	Nothing by mouth after finishing bowel prep
Bowel Prep			START PREP 	FINISH PREP 



Day of Procedure - Outline

- Finish your prep by the designated time on your prep specific instructions.
- Do NOT eat or drink after you complete your prep (including chewing gum). A sip of water with morning medications is OK.
- Bring a driver with you.
- You will be brought back to the preparation area where you will change into a hospital gown and get an IV.
- The GI team will go over the procedure with you and obtain informed consent.
- You will be brought to the endoscopy room, positioned, and given medication for sedation.
- The procedure takes an average of 20-30 minutes.

After your Procedure

- You will be brought to the recovery area.
- You will be informed of the results of your exam.
- You may feel bloated and have some discomfort, but this typically resolves after passing gas.
- For the rest of the day, **NO** strenuous exercise, signing legal documents, operating heavy machinery, driving, or drinking alcohol.
- If you had polyps removed or biopsies taken, you may notice a small amount of blood with the first bowel movement that will later resolve.

Low Residue and Clear Liquid Diet

How to prepare for your colonoscopy

TYPE OF FOOD	LOW RESIDUE DIET [Starting 3 Days Prior]	FOODS TO AVOID
Milk & Dairy	Milk, cheese, buttermilk, cream, yogurt, sour cream	No yogurt mixed with seeds, granola, or berries
Bread & Grains	White bread, white rice, plain crackers, low-fiber cereal (corn flakes, puffed rice)	No brown rice, whole grains, granola, oatmeal, breads with nuts or seeds
Meats & Proteins	Chicken, turkey, lean pork, eggs, tofu, fish, and seafood	No tough meat with gristle, no beans or legumes.
Fruits & Vegetables	Fruit juice without pulp, apple sauce, ripe fruits (cantaloupe, honeydew, peaches), cooked fruits without skin or seeds	No raw fruit with skin or seeds – berries, apples, oranges, watermelon. Raisins, dried fruits.
Nuts & Seeds	Smooth/creamy peanut butter	No seeds (fennel, sesame, pumpkin, sunflower), nuts (peanuts, almonds, walnuts), chunky nut butter, popcorn
Desserts	Ice cream, sorbet, plain pudding, gelatin, cookies or cake made with white flour	No coconut, no whole grain cookies or cakes
Drinks & Beverages	Coffee, tea, hot chocolate, clear fruit drinks (no pulp), soda, sparkling water, Ensure/Boost without added fiber	No beverages with red/purple dyes, no juices with pulp

CLEAR LIQUID DIET [The Entire Day Before Colonoscopy]

- Water
- Clear broth
 - Vegetable, chicken, beef
- Tea or black coffee without cream
- Clear, light-colored juices:
 - Apple
 - White grape
 - Lemonade without pulp
 - White cranberry
- Soda
- Sports beverages
- Flavored water
- Popsicles
- Gelatin



Avoid all red and purple-colored products, as this may look like blood on the day of your procedure!



@EmoryGastroHep
© Emory 2024

SUTAB[®]

Author: Alarica Dietzen, MD

DATE OF
PROCEDURE:

__/__/__

The Day Before Your Procedure

Follow steps 1-5 between 4PM – 6PM then wait AT LEAST 4 HOURS and then repeat steps 1-5.

Take SUTAB at least 1 hour BEFORE or 1 hour AFTER taking your home medications.

Ok to drink: ONLY clear or “see-through” liquids such as yellow Jell-O, chicken broth, water, or Sprite.

After midnight: okay to only have water.

What to expect from your SUTAB prep:

The prep solution can make you feel nauseous or bloated and will induce diarrhea.

Step 1:

Take 12 SUTAB[®] tablets



AND Drink 16 ounces of water within 20 minutes.



Step 2:

After you take the 12th tablet, wait 30 minutes



Step 3:

Drink 16 ounces of water in 30 minutes.



Step 4:

Wait 30 minutes



Step 5:

Drink 16 ounces of water over 30 minutes.



Step 6:

Wait AT LEAST 4 HOURS before repeating STEPS 1-5



The Morning of Your Procedure

4 Hours before your arrival time: After taking your last dose of SUTAB, DO NOT EAT OR DRINK ANYTHING.

At the end of your prep, your stools should be urine-colored!



Call the office if your stools are still formed OR you have not had any bowel movements by 4 hours prior to when you are scheduled to arrive OR if you need anti-nausea medications.